

Monitoring Ionospheric Total Electron Content Using the Global GPS Network

Brian D. Wilson; Anthony J. Mannucci;
Dah-Ning Yuan; Ulf J. Lindqwister; Thomas F. Runge (all at: Jet
Propulsion Laboratory, California Institute of Technology, 4800
Oak Grove Dr., Pasadena, CA 91109; ph. 818-354-2790; e-mail:
bdw@logos.jpl.nasa.gov)

We have developed the capability to monitor the global distribution of ionospheric total electron content (TEC) using dual-frequency observations from a worldwide network of Global Positioning System (GPS) ground receivers. Continuously operating, high-accuracy GPS receivers are currently installed at more than 60 sites around the world covering a wide latitude range (from Ny Alesund, Norway at 78°N to McMurdo, Antarctica at 77°S). By using spatial and temporal interpolation between the TEC measurements, combined with a simple ionospheric shell model, we are able to produce a global snapshot or map of vertical TEC every hour. A time-series of false-color maps can be combined to create an animation which shows the global evolution of ionospheric structure. We can also simultaneously estimate the instrumental biases in the GPS receivers and satellite transmitters, which must be determined in order to extract absolute TEC from the GPS line-of-sight observables.

The global mapping technique utilizes an ionospheric "shell model" approximation in which the ionization is assumed to lie in a thin spherical shell at a fixed height above the Earth, and the line-of-sight measurements are converted to equivalent vertical TEC using an elevation mapping function. Surface interpolation over a triangular grid on the shell is used to combine the individual measurements into a continuous regional or global map. The vertices of the grid are fixed in a solar-geomagnetic coordinate system (geomagnetic latitude and nearly Sun-fixed longitude) since the TEC distribution is slowly varying in that frame. A Kalman-type parameter estimation filter is used to update the ionosphere map as new TEC observations become available. The TEC value at each of 642 vertices is updated once every hour if data near the vertex is available. Otherwise, the vertex value is smoothly interpolated in time between updated values. Kalman filter updating and the use of local grid-based interpolation enable accurate monitoring of sub-hourly changes in the ionosphere over both regional and global scales.

The accuracy of the maps has been assessed by direct comparisons to independent vertical TEC measurements derived from the dual-frequency altimeter on the TOPEX satellite. The TOPEX comparisons indicate that the maps are generally accurate to 5 TECU ($1 \text{ TECU} = 10^{16} \text{ el/m}^2$) in the mid-latitudes and 5-10 TECU in the more variable equatorial region.

American Geophysical Union Abstract Form

Reference # 0000
Session 0.00

1. 1995 Spring Meeting

2. 00 8602343

3. (a) Brian D. Wilson
Mail Stop T1129
Jet Propulsion Laboratory
4800 Oak Grove Dr.
Pasadena, CA 91109 USA
bdw@logos.jpl.nasa.gov

(b) 818-354-2790

(c) 818-393-6890

4. G

5. (a) G07

(b) 2494, 2435, 2479

(c)

6. N/A

7. 20% Radio Sci

8. Charge \$70 to Ulf J. Lindqwister
VISA card 4798 2640 0507 7953,
expires 02/97

9. 1. Program chair: Richard B.
Langley

10. Should be first of several related
papers by J], authors in this
session

11. Regular author

Date received: 8 Feb 95
Date formatted: March 10, 1995
Form version: 1.1